ON THE NON-TRIVIALITY OF THE p-ADIC ABEL-JACOBI IMAGE OF GENERALISED HEEGNER CYCLES MODULO p, I: MODULAR CURVES

ASHAY A. BURUNGALE

arXiv:1410.0300v2 [math.NT] 5 Jun 2015

Abstract. Generalised Heegner cycles are associated with a pair of an elliptic newform and a Hecke character over an imaginary quadratic extension $K/\mathbf{Q}$. Let p be an odd prime split in $K/\mathbf{Q}$ and $l \neq p$ an odd unramified prime. We prove the non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles modulo p over the $\mathbf{Z}_l$-anticylotomic extension of K. The result is an evidence for the refined Bloch-Beilinson and the Bloch-Kato conjecture. In the case of weight two, it provides a refinement of the results of Cornut and Vatsal on Mazur's conjectures regarding the non-triviality of Heegner points over the $\mathbf{Z}_l$-anticylotomic extension of K.

Contents

1. Introduction 1
2. Anticyclotomic toric periods 6
3. Non-triviality of p-adic Abel-Jacobi image modulo p 8
3.1. Non-triviality I 9
3.2. Generalised Heegner cycles 11
3.2.1. Generalities 11
3.2.2. p-adic Abel-Jacobi map 13
3.2.3. Conjectures 14
3.3. Non-triviality II 16
4. p-indivisibility of Heegner points 18
References 19

1. Introduction

One expects the existence of a commutative p-adic L-function $L_p(\mathcal{M})$ associated to a p-ordinary critical motive $\mathcal{M}$ over a tower of number fields with the Galois group being a commutative p-adic Lie-Group Γ, characterised by an interpolation of the p-stabilised critical L-values of $\mathcal{M}$ twisted by a dense subset of characters of Γ. Certain values of the p-adic L-function outside the range of interpolation are expected to encode deep p-adic arithmetic information regarding $\mathcal{M}$. One can ask for the non-triviality of the value of $L_p(\mathcal{M} \otimes \nu)$ at a fixed character outside the range of interpolation, as ν varies over a family of l-power conductor characters, for a prime $l \neq p$. If one has a strategy of approaching the non-triviality of critical L-values of $\mathcal{M}$ twisted by ν as ν varies, it might apply for the former non-triviality as well.

Date: June 8, 2015.
2010 *Mathematics Subject Classification.* Primary 19F27, 11G18, 11R23 Secondary 11F85.

Based on the classical Waldspurger formula (*cf.* [32]), an anticyclotomic p-adic L-function associated with a Rankin-Selberg convolution of an elliptic modular form and a theta series is constructed in [2], [6] and [23]. In [2], values of the p-adic L-function at certain characters outside the range of interpolation are shown to be related to the p-adic Abel-Jacobi image of generalised Heegner cycles associated to the convolution. This can be considered as a p-adic Waldspurger formula (*cf.* [25]). The construction of the cycle is due to Bertolini-Darmon-Prasanna and generalises the one of classical Heegner cycles due to Schoen and Nekovář. The cycle lives in a middle dimensional Chow group of a fiber product of a Kuga-Sato variety arising from a modular curve and a self-product of a CM elliptic curve. In the case of weight two, the cycle coincides with a Heegner point. Based on Hida's strategy, non-triviality of Rankin-Selberg L-values of the l-power conductor anticyclotomic character twists modulo p is shown in [7] and [23]. In this article, we use the strategy to study non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles modulo p over the $\mathbf{Z}_l$-anticyclotomic extension. The non-triviality can be seen as an evidence for the Bloch-Beilinson and the Bloch-Kato conjecture as follows. In the setup, the Rankin-Selberg convolution is self-dual with root number -1. Accordingly, the Bloch-Beilinson conjecture implies the existence of a non-torsion null-homologous cycle in the Chow realisation. In the setup, a natural candidate for a non-trivial null-homologous cycle is the generalised Heegner cycle. For a prime p, the Bloch-Kato conjecture implies the non-triviality of the p-adic étale Abel-Jacobi image of the cycle. A natural question is to further investigate the non-triviality of the p-adic Abel-Jacobi image of the cycle.

In the introduction, for simplicity we mostly restrict to the case of Heegner points.

Let $p > 3$ be an odd prime. We fix two embeddings $\iota_\infty\colon \overline{\mathbf{Q}} \to \mathbf{C}$ and $\iota_p\colon \overline{\mathbf{Q}} \to \mathbf{C}_p$. Let v_p be the p-adic valuation induced by ι_p so that $v_p(p) = 1$. Let $\mathfrak{m}_p$ be the maximal ideal of $\overline{\mathbf{Z}}_p$.

Let $K/\mathbf{Q}$ be an imaginary quadratic extension and $\mathcal{O}$ the ring of integers. Let $-d_K$ be the discriminant. As K is a subfield of the complex numbers, we regard it as a subfield of the algebraic closure $\overline{\mathbf{Q}}$ via the embedding ι_∞. Let G_K be the absolute Galois group $\mathrm{Gal}(\overline{\mathbf{Q}}/K)$. Let c be the complex conjugation on $\mathbf{C}$ which induces the unique non-trivial element of $\mathrm{Gal}(K/\mathbf{Q})$ via ι_∞. We assume the following:

(ord) p splits in K.

Let $\mathfrak{p}$ be a prime above p in K induced by the p-adic embedding ι_p. For an integral ideal $\mathfrak{n}$ of K, we fix a decomposition $\mathfrak{n} = \mathfrak{n}^+\mathfrak{n}^-$ where $\mathfrak{n}^+$ (resp. $\mathfrak{n}^-$) is only divisible by split (resp. ramified or inert) primes in $K/\mathbf{Q}$. For a positive integer m, let H_m be the ring class field of K with conductor m and $\mathcal{O}_m = \mathbf{Z} + m\mathcal{O}$ the corresponding order. Let H be the Hilbert class field.

Let N be a positive integer such that $p \nmid N$. For $k \geq 2$, let $S_k(\Gamma_0(N), \epsilon)$ be the space of elliptic cusp forms of weight k, level $\Gamma_0(N)$ and neben-character ϵ. Let $f \in S_2(\Gamma_0(N), \epsilon)$ be a normalised newform. In particular, it is a Hecke eigenform with respect to all Hecke operators. Let $N_\epsilon | N$ be the conductor of ϵ. Let E_f be the Hecke field of f and $\mathcal{O}_{E_f}$ the ring of integers. Let $\mathfrak{P}$ be a prime above p in E_f induced by the p-adic embedding ι_p. Let $\rho_f : \mathrm{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \to \mathrm{GL}_2(\mathcal{O}_{E_f,\mathfrak{P}})$ be the corresponding p-adic Galois representation.

We assume the following generalised Heegner hypothesis:

(Hg) $\mathcal{O}$ contains an ideal $\mathfrak{N}$ of norm N such that there exists an isomorphism

$$\mathcal{O}/\mathfrak{N} \simeq \mathbf{Z}/N\mathbf{Z}.$$

From now, we fix such an ideal $\mathfrak{N}$. Let $\mathfrak{N}_\epsilon|\mathfrak{N}$ be the unique ideal of norm N_ϵ.

Let $\mathbf{N}$ denote the norm Hecke character over $\mathbf{Q}$ and $\mathbf{N}_K := \mathbf{N} \circ N_{\mathbf{Q}}^K$ the norm Hecke character over K. For a Hecke character λ over K, let $\mathfrak{f}_\lambda$ (resp. ϵ_λ) denote its conductor (resp. the restriction $\lambda|_{\mathbf{A}_{\mathbf{Q}}^\times}$, where $\mathbf{A}_{\mathbf{Q}}$ denotes the adele ring over $\mathbf{Q}$). We say that λ is central critical for f if it is of infinity type (j_1, j_2) with $j_1 + j_2 = 2$ and $\epsilon_\lambda = \epsilon\mathbf{N}^{\mathbf{2}}$.

Let b be a positive integer prime to pN. Let $\Sigma_{cc}(b, \mathfrak{N}, \epsilon)$ be the set of Hecke characters λ such that:

(C1) λ is central critical for f,
(C2) $\mathfrak{f}_\lambda = b \cdot \mathfrak{N}_\epsilon$ and
(C3) The local root number $\epsilon_q(f, \lambda^{-1}) = 1$, for all finite primes q.

Let χ be a finite order Hecke character such that $\chi\mathbf{N}_K \in \Sigma_{cc}(b, \mathfrak{N}, \epsilon)$. Let $E_{f,\chi}$ be the finite extension of E_f obtained by adjoining the values of χ.

Let $X_1(N)$ be the modular curve of level $\Gamma_1(N)$, ∞ the standard cusp $i\infty$ of $X_1(N)$ and $J_1(N)$ the corresponding Jacobian. Let B_f be the abelian variety associated to f by the Eichler-Shimura correspondence and $\Phi_f : J_1(N) \to B_f$ the associated surjective morphism. The uniqueness of the abelian variety follows from the assumption that f is a newform. By possibly replacing B_f with an isogenous abelian variety, we suppose that B_f endomorphisms by the integer ring $\mathcal{O}_{E_f}$. Let ω_f be the differential form on $X_1(N)$ corresponding to f. We use the same notation for the corresponding 1-form on $J_1(N)$. Let $\omega_{B_f} \in \Omega^1(B_f/E_f)^{\mathcal{O}_{E_f}}$ be the unique 1-form such that $\Phi_f^*(\omega_{B_f}) = \omega_f$. Here $\Omega^1(B_f/E_f)^{\mathcal{O}_{E_f}}$ denotes the subspace of 1-forms given by

$$\Omega^1(B_f/E_f)^{\mathcal{O}_{E_f}} = \left\{\omega \in \Omega^1(B_f/E_f) | [\lambda]^*\omega = \lambda\omega, \forall \lambda \in \mathcal{O}_{E_f}\right\}.$$

Recall that b is a positive integer prime to N. Let A_b be an elliptic curve with endomorphism ring $\mathcal{O}_b = \mathbf{Z} + b\mathcal{O}$, defined over the ring class field H_b. Let t be a generator of $A_b[\mathfrak{N}]$. We thus obtain a point $x_b = (A_b, A_b[\mathfrak{N}], t) \in X_1(N)(H_{bN})$. Let $\Delta_b = [A_b, A_b[\mathfrak{N}], t] - (\infty) \in J_1(N)(H_{bN})$ be the corresponding Heegner point on the modular Jacobian. We regard χ as a character $\chi : \mathrm{Gal}(H_{bN}/K) \to E_{f,\chi}$. Let H_χ be the abelian extension of K cut out by the character χ. To the pair (f, χ), we associate the Heegner point $P_f(\chi)$ given by

$$P_f(\chi) = \sum_{\sigma \in \mathrm{Gal}(H_{bN}/K)} \chi^{-1}(\sigma)\Phi_f(\Delta_b^\sigma) \in B_f(H_\chi) \otimes_{\mathcal{O}_{E_f}} E_{f,\chi}. \tag{1.1}$$

To consider the non-triviality of the Heegner points $P_f(\chi)$ as χ varies, we can consider the non-triviality of the corresponding p-adic formal group logarithm. The restriction of the p-adic logarithm gives a homomorphism

$$\log_{\omega_{B_f}} : B_f(H_\chi) \to \mathbf{C}_p.$$

We extend it to $B_f(H_\chi) \otimes_{\mathcal{O}_{E_f}} E_{f,\chi}$ by $E_{f,\chi}$-linearity.

We now fix a finite order Hecke character η such that $\eta\mathbf{N}_K \in \Sigma_{cc}(c, \mathfrak{N}, \epsilon)$, for some c. For $v|c^-$, let $\Delta_{\eta,v}$ be the finite group $\eta(\mathcal{O}_{K_v}^\times)$. Here $\mathcal{O}_{K_v}$ denotes the integer ring of the local field K_v. Let

$l \neq p$ be an odd prime unramified in K and prime to cN. Let $H_{cNl^\infty} = \bigcup_{n\geq 0} H_{cNl^n}$ be the ring class field of conductor cNl^∞. Let $G_n = \mathrm{Gal}(K_{cNl^n}/K)$ and $\Gamma_l = \varprojlim G_n$. Let $\mathfrak{X}_l$ denote the set of l-power order anticyclotomic characters of Γ_l. As $\nu \in \mathfrak{X}_l$ varies, we consider the non-triviality of $\log_{\omega_{B_f}}(P_f(\eta\nu))/p$ modulo p.

Our result is the following.

Theorem A. Let $f \in S_2(\Gamma_0(N), \epsilon)$ be a normalised newform and η a finite order Hecke character such that $\eta\mathbf{N}_K \in \Sigma_{cc}(c, \mathfrak{N}, \epsilon)$, for some c. In addition to the hypotheses (ord) and (Hg), suppose that

(1). The residual representation $\rho_f|_{G_K}$ mod $\mathfrak{m}_p$ is absolutely irreducible,
(2). $\mathfrak{N}^-$ is square-free and
(3). $p \nmid \prod_{v|c^-} \Delta_{\eta,v}$.

Then, for all but finitely many $\nu \in \mathfrak{X}_l$ we have

$$v_p\left(\frac{\log_{\omega_{B_f}}(P_f(\eta\nu))}{p}\right) = 0.$$

We have the following immediate corollary.

Corollary A. Let $f \in S_2(\Gamma_0(N), \epsilon)$ be a normalised newform and η a finite order Hecke character such that $\eta\mathbf{N}_K \in \Sigma_{cc}(c, \mathfrak{N}, \epsilon)$, for some c. In addition to (Hg), suppose that

(1). f does not have CM over K and
(2). $\mathfrak{N}^-$ is square-free.

Then, for all but finitely many $\nu \in \mathfrak{X}_l$ the Heegner points $P_f(\eta\nu)$ are non-zero in $B_f(H_{\eta\nu}) \otimes_{T_f} E_{f,\eta\nu}$.

Proof. This follows from the existence of a prime p satisfying the hypotheses of Theorem A. □

For analogous non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles modulo p, we refer to §3.3.

We now describe strategy of the proof. Some of the notation used here is not followed in the rest of the article. We basically follow a strategy of Hida. To slightly strengthen the strategy, we first define anticyclotomic toric periods $P_{g,\lambda}(\nu, n)$ associated to a pair (g, λ). Here g is a p-adic elliptic modular form, λ a Hecke character and ν is as above factoring through G_n. We then state a non-triviality theorem implying the non-triviality of the toric periods modulo p as long as g is a p-integral nearly holomorphic elliptic modular form defined over a number field and we have non-triviality of a certain

toric form associated to the pair modulo p. The theorem follows from Hida's proof of the non-triviality of an anticyclotomic modular measure modulo p in [17]. The later is based on Chai's theory of Hecke stable subvarieties of a mod p Shimura variety (*cf.* [15]). Let d be the Katz p-adic differential operator and $f^{(p)}$ the p-depletion of f. For the pair $(d^{-1}(f^{(p)}), \eta)$, the toric period $P_{d^{-1}(f^{(p)}),\eta}(\nu, n)$ essentially equals $\log_{\omega_{B_f}}(P_f(\eta\nu))/p$. This follows from the p-adic Waldspurger formula. As $d^{-1}(f^{(p)})$ is a weight zero p-adic elliptic modular form, the non-triviality theorem does not directly apply. However, we show that there exists a pair (g, λ) such that its toric periods are congruent modulo p to the toric periods of the interest and the non-triviality theorem applies. The nearly holomorphic elliptic modular form g is an iterated image under the p-adic differential operator of f and the congruence is a rather simple consequence of the q-expansion principle. The criterion in the non-triviality theorem is basically verified for the pair (g, λ) in [23] based on the computation of Fourier coefficients of a toric form. The hypotheses (1) and (2) guarantee the non-triviality of the toric form modulo p.

We would like to remark that neither L-values nor p-adic L-functions appear explicitly in the statement or the proof of Theorem A. The above strategy can be also used to prove the non-triviality of the p-adic formal group logarithm arising from toric forms in [23] of the Heegner points on the modular Jacobian modulo p .

A related modulo p non-triviality is also considered in [34]. As [loc. cit.] treats the case when p is inert in K, our result is complementary. Without the hypotheses (1) and (2), Corollary A was conjectured by Mazur (*cf.* [26]). Under the classical Heegner hypotheses, it was indepdendently proven by Cornut and Vatsal (*cf.* [11], [33], [34] and also [12]). For a related work of Aflalo- Nekovář, we refer to [1]. In [11], there is an additional hypothesis that the corresponding abelian variety B_f has good ordinary reduction at p. As far as we know, Corollary A is a first result regarding Mazur's conjecture under the generalised Heegner hypotheses (Hg). Moreover, the theorem is a first result regarding the non-triviality of the p-adic formal group logarithm of Heegner points modulo p. Our approach is new. It seems suggestive to compare our approach with the earlier approaches. Here we only mention the following and for more refer to [9]. In [34], Jochnowitz congruence is a starting point. It reduces the non-triviality of the Heegner points to the non-triviality of the Gross points on a suitable definite Shimura "variety". This in turn is studied via Ratner's theorem. In this article, the non-triviality is based on the modular curve itself. As indicated above, it relies on Chai's theory of Hecke stable subvarieties of a mod p Shimura variety. In other words, the result of Cornut and Vatsal now fits in the framework of Hida (*cf.* [17]). An expectation along these lines was expressed in [35]. Before the p-adic Waldspurger formula, the result and the framework appeared to be complementary. The formula also allows a rather smooth transition to the higher weight case.

As far as we know, higher weight analogue of Theorem A is a first general result regarding the non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles. The only known result seems to be the non-triviality of several examples of such families in [4]. Howard proved an analogous non-triviality of the étale Abel-Jacobi image of classical Heegner cycles in [21].

Theorem A and its higher weight analogue has various arithmetic applications besides the Bloch-Beilinson and the Bloch-Kato conjecture. Consequences for the p-indivisibility of Heegner points when f is p-ordinary and the refined Bloch-Beilinson conjecture are described in §4 and the remark (1) following Theorem 3.5, respectively. The later is based on [4]. The non-triviality also implies that the Griffiths group of the fiber product of a Kuga-Sato variety arising from the Modular curve and a self-product of a CM elliptic curve has infinite rank over the anticyclotomic extension. Applications to the infinitude of the Griffiths groups modulo p will be considered in the near future. Arithmetic

consequences of Corollary A are well documented in the literature, for example [11] and [34].

In [8], we prove an analogue of the results for Shimura curves over the rationals in the case $l = p$. Recently, the p-adic Waldspurger formula has been generalised to modular forms on Shimura curves over a totally real field (*cf.* [25]). In the future, we hope to consider an analogous non-triviality of generalised Heegner cycles over a CM field.

The article is perhaps a follow up to the articles [2] and [17]. We refer to them for a general introduction. For a survey, we refer to [9].

The article is organised as follows. In §2, we describe a few generalities regarding the anticyclotomic toric periods. In §3, we prove non-triviality of generalised Heegner cycles modulo p. For simplicity, we first treat the case of Heegner points. In §3.1, we first state the p-adic Waldspurger formula and then prove Theorem A. In §3.2, we describe generalities regarding generalised Heegner cycles and state conjectures regarding their non-triviality. In §3.3, we prove the non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles modulo p. In §4, we prove the p-indivisibility of Heegner points when f is p-ordinary.

Acknowledgments. We are grateful to our advisor Haruzo Hida for continuous guidance and encouragement. We thank Ming-Lun Hsieh for many instructive comments on the previous versions of the article and also for his assistance. We thank Francesc Castella for helpful conversations, particularly regarding the p-adic Waldspurger formula. We thank Henri Darmon and Burt Totaro for instructive comments. Finally, we thank Miljan Brakocevic, Brian Conrad, Samit Dasgupta, Chandrashekhar Khare, Jan Nekovář, Vinayak Vatsal, Kevin Ventullo and Xinyi Yuan for helpful conversations about the topic.

Notation We use the following notation unless otherwise stated.

For a number field L, let $\mathcal{O}_L$ be the ring of integers, $\mathbf{A}_L$ the adele ring, $\mathbf{A}_{L,f}$ the finite adeles and $\mathbf{A}_{L,f}^{(\square)}$ the finite adeles away from a finite set of places $\square$ of L. For a fractional ideal $\mathfrak{a}$, let $\widehat{\mathfrak{a}} = \mathfrak{a} \otimes \widehat{\mathbf{Z}}$. Let G_L be the absolute Galois group of L and G_L^{ab} the maximal abelian quotient. Let $\mathrm{rec}_L : \mathbf{A}_L^\times \to G_L^{ab}$ be the geometrically normalized reciprocity law.

We refer to an elliptic modular form as a modular form. For a modular form g, let its Fourier expansion $g(q)$ at a cusp $\mathbf{c}$ be given by

$$g(q) = \sum_{n \geq 0} \mathbf{a}_n(g, \mathbf{c}) q^n.$$

2. Anticyclotomic toric periods

In this section, we describe a few generalities regarding anticyclotomic toric periods and their non-triviality modulo p. The section is based on Hida's approach in [17] and [19], as formulated in [22]. We give a slightly different formulation and a mild strengthening.

Let the notations and hypotheses be as in the introduction. For a non-negative integer n, recall that H_{cNl^n} is the ring class field of K of conductor cNl^n. Let $R_n = \mathbf{Z} + cNl^n\mathcal{O}$. We have an identification $\mathrm{Pic}(R_n) = G_n$ such that $a\widehat{R}_n \cap K \leftrightarrow \mathrm{rec}_K(a)|_{H_{cNl^n}}$. Let $U_n = \mathbf{C}_1 \times (\widehat{R_n})^\times \subset \mathbf{C}^\times \times \mathbf{A}_{K,f}^\times$ be a compact subgroup, where $\mathbf{C}_1$ is the unit complex circle. Via the reciprocity law, we have the identification $G_n = K^\times \mathbf{A}_{\mathbf{Q}}^\times \backslash \mathbf{A}_K^\times / U_n$. Let $[\cdot]_n : \mathbf{A}_K^\times \to G_n$ be the quotient map and $[a] = \varprojlim_n [a]_n \in \Gamma_l$. For $a \in \mathbf{A}_K^\times$, let $x_n(a)$ be the CM point $x([a]_n)$ in [22, §3.3]. In the notation of the introduction, $x(\sigma)$ is the CM point $x_{cl^n}^\sigma$ along with a choice of p^∞-level structure. Let $\mathbf{c}(a)$ be the polarisation ideal of the CM point $x_0(a)$ in [22, §3.4].

Let $K_0 \subset \mathrm{GL}_2(\mathbf{A}_{\mathbf{Q},f})$ be an open compact subgroup and let $S_k(K_0, A)$ be the space of cusp forms of weight k and level K_0 over A, where A is $\mathbf{Z}_{(p)}$-algebra. Let $V(K_0, O)$ be the space of p-adic modular forms of level K over O, where O is a finite flat $\mathbf{Z}_p$-algebra. For the geometric definitions, we refer to [18, §4.2].

Let λ be an arithmetic Hecke character and $\widehat{\lambda}$ its p-adic avatar. We recall that $\widehat{\lambda} : \mathbf{A}_{K,f}^\times / K^\times \to \mathbf{C}_p^\times$ is given by

$$\widehat{\lambda}(x) = \lambda(x) x_p^{(j+\kappa_c(1-c))}.$$

Here $x \in \mathbf{A}_{K,f}^\times$ and λ is of infinity type $(j + \kappa_c, \kappa_c)$.

A p-adic modular form is a formal function on the Igusa tower (*cf.* [20, 7.2.4]). In particular, it is a function of an isomorphism class of a pair consisting of a p-ordinary point on the underlying modular curve and a p^∞-level structure on the point.

Let $g \in V(K_0, O)$ such that the following condition holds:

(T) $g(x_n(ta)) = \widehat{\lambda}(a)^{-1} g(x_n(t))$, where $a \in U_n \mathbf{A}_{\mathbf{Q}}^\times$.

We now define anticyclotomic toric periods of g (compare [17, §3.1 and §3.4]). For $\phi : G_n \to \overline{\mathbf{Z}}_p$, let $P_{g,\lambda}(\phi, n)$ be the $\overline{\mathbf{Z}}_p$-valued toric period given by

$$P_{g,\lambda}(\phi, n) = \sum_{[t]_n \in G_n} \phi([t]_n) g(x_n(t)) \widehat{\lambda}(t). \tag{2.1}$$

In view of (T), the above expression is well defined. We remark that the toric period typically depends on n and in general, the above definition may not give rise to a measure on Γ_l. Under additional hypothesis on g, we can normalise the toric periods and thereby obtain a measure on Γ_l (*cf.* [17, §3.1 and §3.4]).

For the rest of the section, we now suppose that g is a p-integral nearly holomorphic modular form defined over a number field.

To discuss the non-triviality of the toric periods modulo p, we introduce more notation. Let Δ_l be the torsion subgroup of Γ_l, Γ_l^{alg} the subgroup of Γ_l generated by $[a]$ for $a \in (\mathbf{A}_K^{(lp)})^\times$ and $\Delta_l^{alg} = \Gamma_l^{alg} \cap \Delta_l$. Let $\mathcal{R} = \{r\}$ be a set of representatives of Δ_l^{alg} in $(\mathbf{A}_K^{(lp)})^\times$. Let $\rho : \mathbf{A}_K^\times \hookrightarrow \mathrm{GL}_2(\mathbf{A}_{\mathbf{Q}})$ be the toric embedding and $\varsigma \in \mathrm{GL}_2(\mathbf{A}_K^{(lp)})$ as in [22, §3.1 and §3.2], respectively. For $a \in (\mathbf{A}_K^{(lp)})^\times$, let

$$g|[a] = g|\rho_\varsigma(a), \rho_\varsigma(a) = \varsigma^{-1}\rho(a)\varsigma. \tag{2.2}$$

For the definition of $|[a]$ operator, we refer to [22, §2.6].

Let

$$g^{\mathcal{R}} = \sum_{r \in \mathcal{R}} \widehat{\lambda}(r) g|[r]. \tag{2.3}$$

We consider the following hypothesis.

(H) For every $u \in \mathbf{Z}$ prime to l and a positive integer r, there exists a positive integer $\beta \equiv u \bmod l^r$ and $a \in \mathbf{A}_{K,f}^{\times}$ such that $\mathbf{a}_{\beta}(g^{\mathcal{R}}, \mathbf{c}(a)) \neq 0 \bmod \mathfrak{m}_p$.

We have the following result regarding the non-triviality of the toric periods modulo p.

Theorem 2.1. *(Hida) Let the notations be as above. In addition to (ord), suppose that the hypothesis (H) holds. Then, we have*

$$P_{g,\lambda}(\nu, n) \neq 0 \, mod \, \mathfrak{m}_p,$$

for all but finitely many $\nu \in \mathfrak{X}_l$ factoring through G_n (as $n \to \infty$, cf. [17, Thm. 3.2 and Thm 3.3]*).*

Proof. This follows from the proof of the non-triviality of the anticyclotomic modular measure modulo p in [17]. The additional hypothesis that the modular form g is an U_l-eigenform is only used in the construction of the measure. The proof of the non-triviality only relies on the vanishing of the toric periods modulo p and the hypothesis (H).

□

Remark. (1). As the theorem does not require the nearly holomorphic form to be an U_l-eigenform, it can be used to mildly simplify the proof of (l, p)-non-triviality of L-values in [22] and [23].

(2). The Chai-Oort rigidity principle predicts that a Hecke stable subvariety of a mod p Shimura variety is a Shimura subvariety. The principle has been studied by Chai in a series of articles (*cf.* [13], [14] and [15]). The theorem crucially relies on the principle.

3. Non-triviality of p-adic Abel-Jacobi image modulo p

In this section, we prove the non-triviality of generalised Heegner cycles modulo p. For simplicity, we first treat the case of Heegner points. In §3.1, we first state the p-adic Waldspurger formula and then prove the non-triviality of the p-adic formal group logarithm of Heegner points modulo p (*cf.* Theorem A). In §3.2, we describe generalities regarding generalised Heegner cycles and state conjectures regarding their non-triviality. In §3.3, we prove the non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles modulo p.

3.1. **Non-triviality I.** In this subsection, we first state the p-adic Waldspurger formula and then prove the non-triviality of the p-adic formal group logarithm of Heegner points modulo p (*cf.* Theorem A).

Let the notations and hypotheses be as in the introduction. In particular, f is a weight two Hecke-eigen cuspform and χ a finite order Hecke character such that $\chi\mathbf{N}_K \in \Sigma_{cc}(b, \mathfrak{N}, \epsilon)$.

Let V and U be the Hecke operators on the space $V(\Gamma_0(N), O)$ as defined in [2, §3.8]. For $g \in V(\Gamma_0(N), O)$, let $g^{(p)}$ be the p-depletion given by

$$g^{(p)} = g|(VU - UV). \tag{3.1}$$

The Fourier expansion of $g^{(p)}$ at a cusp $\mathbf{c}$ is given by

$$g^{(p)}(q) = \sum_{p\nmid n} \mathbf{a}_n(g, \mathbf{c})q^n \tag{3.2}$$

(*cf.* [2, (3.8.4)]).

Recall that d denotes the Katz p-adic differential operator (*cf.* [24, Ch. II] and [20, Ch. I]). In view of (3.2), it follows that $d^{-1}(g^{(p)}) = \lim_{j\to 0} d^{-1+j}(g^{(p)})$ is a weight zero p-adic modular forms.

We have the following formula for the anticyclotomic toric periods of the weight zero p-adic modular form $d^{-1}(f^{(p)})$.

Theorem 3.1. *(Bertolini-Darmon-Prasanna) Under the notations and hypotheses as above, we have*

$$\sum_{\sigma\in G_b} \chi^{-1}(\sigma)d^{-1}(f^{(p)})(x(\sigma)) = (1 - \chi^{-1}(\overline{\mathfrak{p}})p^{-1}\mathbf{a}_p(f) + \chi^{-2}(\overline{\mathfrak{p}})\epsilon(p)p^{-1}) \log_{\omega_{B_f}}(P_f(\chi)).$$

Proof. This is proven in [2] and [3]. As the result is stated there in terms of a Rankin-Selberg p-adic L-function, we give some details.

We first recall that

$$T_p(f) = \mathbf{a}_p(f)f, \langle p\rangle(f) = \epsilon(p)f. \tag{3.3}$$

Here $\langle p\rangle$ denotes the diamond operator associated with p.

Let F_f the Coleman primitive of the differential ω_f associated with f (*cf.* [2, §3.7 and §3.8]). In view of (3.3) and [2, Lem. 3.23 and Prop. 3.24], it follows that

$$F_f(x(\sigma)) = d^{-1}(f^{(p)})(x(\sigma)) - \epsilon(p)\mathbf{a}_p(f)d^{-1}(f^{(p)})(\mathfrak{p} * x(\sigma)) + \frac{\epsilon(p)}{p}d^{-1}(f^{(p)})(\mathfrak{p}^2 * x(\sigma)). \tag{3.4}$$

From [2, Lem. 3.23] and [3, Thm. 3.12], we have

$$F_f(x(\sigma)) = \log_{\omega_{B_f}}(\Phi_f(\Delta^\sigma)). \tag{3.5}$$

Here the "$*$"-action on the CM point is as in [2, (1.4.8)].

The summation of the above identity over $\sigma \in G_b$ along with (3.5) and [2, Proof of Thm. 5.13]

finishes the proof.

□

We now consider the variation of the above identity over the $\mathbf{Z}_l$-anticyclotomic extension of K. We fix a finite order Hecke character η such that $\eta\mathbf{N}_K \in \Sigma_{cc}(c, \mathfrak{N}, \epsilon)$, for some c. In what follows, we apply the formalism in §2 to the pair $(d^{-1}(f^{(p)}), \eta)$.

As $d^{-1}(f^{(p)})$ is a weight zero p-adic modular form, the hypothesis (T) holds and the toric periods are well defined. As η is of finite order with conductor $c\mathfrak{N}_\epsilon$, for $\phi : G_n \to \overline{\mathbf{Z}}_p$ we have

$$P_{d^{-1}(f^{(p)}),\eta}(\phi, n) = \sum_{[t]_n \in G_n} (\phi\eta)([t]_n) d^{-1}(f^{(p)})(x_n(t)). \tag{3.6}$$

We have the following formula for the toric periods.

Lemma 3.2. *For $\nu \in \mathfrak{X}_l$, we have*

$$P_{d^{-1}(f^{(p)}),\eta}(\nu, n) = (1 - (\eta\nu)^{-1}(\overline{\mathfrak{p}})p^{-1}\mathbf{a}_p(f) + (\eta\nu)^{-2}(\overline{\mathfrak{p}})\epsilon(p)p^{-1}) \log_{\omega_{B_f}}(P_f(\eta\nu)). \tag{3.7}$$

Proof. We first note that as $\eta\mathbf{N}_K \in \Sigma_{cc}(c, \mathfrak{N}, \epsilon)$, it follows that $\eta\nu\mathbf{N}_K \in \Sigma_{cc}(cl^k, \mathfrak{N}, \epsilon)$ for a non-negative integer k. The lemma thus immediately follows from the previous theorem.

□

We are now ready to prove Theorem A (*cf.* §1).

Theorem 3.3. *Let $f \in S_2(\Gamma_0(N), \epsilon)$ be a normalised newform and η a finite order Hecke character such that $\eta\mathbf{N}_K \in \Sigma_{cc}(c, \mathfrak{N}, \epsilon)$, for some c. In addition to the hypotheses (ord) and (Hg), suppose that*

(1). The residual representation $\rho_f|_{G_K}$ mod $\mathfrak{m}_p$ is absolutely irreducible,
(2). $\mathfrak{N}^-$ is square-free and
(3). $p \nmid \prod_{v|c^-} \Delta_{\eta,v}$.

Then, for all but finitely many $\nu \in \mathfrak{X}_l$ we have

$$v_p\left(\frac{\log_{\omega_{B_f}}(P_f(\eta\nu))}{p}\right) = 0.$$

Proof. In view of Lemma 3.2, it suffices to prove the non-triviality of the toric periods modulo p. However, Theorem 2.1 cannot be directly applied as $d^{-1}(f^{(p)})$ is not a nearly holomorphic modular form.

In view of the effect of the Katz p-adic differential operator on the q-expansion and the q-expansion principle, we have

$$d^{(p-2)p}(f^{(p)}) \equiv d^{-1}(f^{(p)}) \bmod \mathfrak{m}_p$$

and

$$d^{(p-2)p}(f^{(p)}) \equiv d^{(p-2)p}(f) \bmod \mathfrak{m}_p.$$

It thus follows that

$$\sum_{[t]_n\in\Gamma_n}\phi([t]_n)d^{(p-2)p}(f)(x_n(t))(\mathbf{N}_K^{(p-1)^2}\eta)(t)$$
$$\equiv\sum_{[t]_n\in\Gamma_n}\phi([t]_n)d^{-1}(f^{(p)})(x_n(t))\eta(t)\bmod\mathfrak{m}_p.$$

As $d^{(p-2)p}(f)$ is nearly holomorphic of weight $2(p-1)^2$, the pair $(d^{(p-2)p}(f),\eta\mathbf{N}_K^{(p-1)^2})$ satisfies (T) and the toric periods are well defined. The above congruence can now be rewritten as

$$P_{d^{(p-2)p}(f),\mathbf{N}_K^{(p-1)^2}\eta}(\phi,n)\equiv P_{d^{-1}(f^{(p)}),\eta}(\phi,n)\bmod\mathfrak{m}_p. \tag{3.8}$$

Now, we can and do apply Theorem 2.1 to the pair $(d^{(p-2)p}(f),\eta\mathbf{N}_K^{(p-1)^2})$. It suffices to verify the following hypothesis.

(H') For every $u\in\mathbf{Z}$ prime to l and a positive integer r, there exists a positive integer $\beta\equiv u\bmod l^r$ and $a\in\mathbf{A}_{K,f}^{\times}$ such that $v_p(\mathbf{a}_\beta(d^{(p-2)p}(f^{\mathcal{R}}),\mathbf{c}(a)))=0$.

Under the hypotheses (1)-(3), this is essentially verified in [23, §7.4]. We give a brief summary. In view of [23, Lem. 7.2] and [22, Rem. 6.2], it suffices to verify (H') for the toric form $\varphi^{\circ}_{\eta\mathbf{N}_K^{(p-1)^2}}$ in [23, Rem. 3.15]. Based on the computation of local Fourier coefficients of the toric form in [23, §3.10], the hypothesis (H') can be verified as in the proof of [23, Thm. 6.2]. We refer to the proof of [23, Thm. 6.2] for related details.

In view of the congruence (3.8), this finishes the proof.

As the non-triviality of the p-adic formal group logarithm of a point implies the point being non-torsion, "In particular" part of the Theorem follows.

□

Remark. The hypothesis on p being odd arises from the corresponding hypothesis in [17]. The hypotheses (1)-(3) intervene only in the computation of the Fourier coefficients of the toric form. It is expected that the hypothesis (2) could be removed. However, the hypotheses (1) and (3) are perhaps essential. We refer to the introduction of [23] for a related discussion.

3.2. **Generalised Heegner cycles.** In this section, we describe basic setup regarding generalised Heegner cycles.

3.2.1. *Generalities.* We briefly recall generalities regarding generalised Heegner cycles following [2, §2] and [4, §4].

Unless otherwise stated, let the notations and hypotheses be as in the introduction.

Recall that $X_1(N)$ is the modular curve of level $\Gamma_1(N)$ and ∞ the standard cusp $i\infty$. Here i is a chosen square root of -1. Strictly speaking, the modular curve only exists as a Deligne-Mumford stack for $N \leq 3$. Let r be a non-negative integer. Let W_r be the r-fold Kuga-Sato variety over $X_1(N)$ constructed in [2, App.]. In other words, W_r is the canonical desingularisation of the r-fold self-product of the universal elliptic curve over $X_1(N)$.

Let A be a CM elliptic curve with CM by $\mathcal{O}$ defined over the Hilbert class field H (*cf.* [2, §1.4]). Let d be a positive integer prime to N. For an ideal class $[\mathfrak{a}] \in \mathrm{Pic}(\mathcal{O}_{dN})$ with $\mathfrak{a}$ prime to N, let $\varphi_{\mathfrak{a}}$ be the natural isogeny

$$\varphi_{\mathfrak{a}} : A \to A_{\mathfrak{a}} = A/A[\mathfrak{a}]. \tag{3.9}$$

Strictly speaking, the above definition of the CM elliptic curve $A_{\mathfrak{a}}$ is correct only when $d = 1$ and a suitable $\mathcal{O}$-transform needs to be considered in the general case (*cf.* [2, §1.4]).

Let r_1 and r_2 be non-negative integers such that $r_1 \geq r_2$ and $r_1 \equiv r_2 \bmod 2$. Let s and u be non-negative integers such that

$$r_1 + r_2 = 2s \text{ and } r_1 - r_2 = 2u.$$

Let X_{r_1,r_2} be a $r_1 + r_2 + 1$-dimensional variety given by

$$X_{r_1,r_2} = W_{r_1} \times A^{r_2}.$$

For an ideal $\mathfrak{a}$ as above, we consider

$$(A_{\mathfrak{a}} \times A)^{r_2} \times (A_{\mathfrak{a}} \times A_{\mathfrak{a}})^u = (A_{\mathfrak{a}}^{r_1} \times A^{r_2}) \subset X_{r_1,r_2}.$$

The last inclusion arises from the embedding of $A_{\mathfrak{a}}^{r_1}$ in W_{r_1} as a fiber of the natural projection $W_{r_1} \to X_1(N)$.

Let $\Gamma_{\mathfrak{a}} \in Z^1(A_{\mathfrak{a}} \times A_{\mathfrak{a}})$ be the transpose of the graph of $\sqrt{-d_K}$. Here $\sqrt{-d_K}$ is the square root of $-d_K$ whose image under the complex embedding ι_∞ has positive imaginary part. Let $\Gamma_{\varphi,\mathfrak{a}} \in Z^1(A_{\mathfrak{a}} \times A)$ be the transpose of the graph of $\varphi_{\mathfrak{a}}$. Let

$$\Gamma_{r_1,r_2,\mathfrak{a}} = \Gamma_{\varphi,\mathfrak{a}}^{r_2} \times \Gamma_{\mathfrak{a}}^{u}$$

and

$$\Delta_{r_1,r_2,\mathfrak{a}} = \epsilon_{X_{r_1,r_2}}(\Gamma_{r_1,r_2,\mathfrak{a}}) \in CH^{s+1}(X_{r_1,r_2} \otimes L)_{0,\mathbf{Q}}. \tag{3.10}$$

Here $\epsilon_{X_{r_1,r_2}}$ is the idempotent in the ring of correspondences on X_{r_1,r_2} defined in [4, §4.1] and L is the field of definition of the cycle $\Gamma_{r_1,r_2,\mathfrak{a}}$. The idempotent has the effect of making the cycle null-homologous (*cf.* [4, Prop. 4.1.1]). The notation $CH^{s+1}(X_{r_1,r_2} \otimes L)_{0,\mathbf{Q}}$ denotes the Chow group of homologically trivial cycles over L of codimension $s+1$ with rational coefficients.

When $r_1 = r_2 = r$, we let X_r, $\Gamma_{r,\mathfrak{a}}$ and $\Delta_{r,\mathfrak{a}}$ denote $X_{r,r}$, $\Gamma_{r,r,\mathfrak{a}}$ and $\Delta_{r,r,\mathfrak{a}}$, respectively.

When r_1 is even and $r_2 = 0$, the above cycles are nothing but the classical Heegner cycles (*cf.* [2, §2.4]). Classical Heegner cycles were first investigated in [31] and [28].

When $r = 0$, the generalised Heegner cycle $\Delta_{0,\mathfrak{a}}$ is a CM point on the modular curve $X_1(N)$. In this case, we replace $\Delta_{0,\mathfrak{a}}$ by $\Delta_{0,\mathfrak{a}} - \infty$ to make it homologically trivial. Note that this preserves the field of definition of the cycles as the ∞-cusp is defined over $\mathbf{Q}$.

3.2.2. *p-adic Abel-Jacobi map.* We briefly recall generalities regarding a relevant p-adic Abel-Jacobi map following [2, §3]. For the dual exponential map, we also refer to [5].

Let the notations and hypotheses be as in §3.2.1. Let ϵ_X denote the idempotent $\epsilon_{X_{r_1,r_2}}$.

Let F be a number field containing the Hilbert class field H. Let V_s be the p-adic Galois representation of G_F given by

$$V_s = H^{2s+1}_{\acute{e}t}(X_{r_1,r_2} \times_F \overline{\mathbf{Q}}, \mathbf{Q}_p).$$

Let $AJ^{\acute{e}t}_F$ be the étale Abel-Jacobi map

$$AJ^{\acute{e}t}_F : CH^{s+1}(X_{r_1,r_2} \otimes F)_{0,\mathbf{Q}} \to H^1(F, \epsilon_X V_s(s+1)) \tag{3.11}$$

due to S. Bloch (*cf.* [28] and [2, Def. 3.1]). The Bloch-Kato conjecture implies that the $\mathbf{Q}_p$-linearisation $AJ^{\acute{e}t}_F \otimes \mathbf{Q}_p$ is injective (*cf.* [28, (2.1)]).

Let v be a place in F above p induced by the p-adic embedding ι_p. We have the localisation map

$$loc_v : H^1(F, \epsilon_X V_s(s+1)) \to H^1(F_v, \epsilon_X V_s(s+1))$$

in Galois cohomology.

In general, the localisation map need not be injective.

The image of the composition $loc_v \circ AJ^{\acute{e}t}_F$ is contained in the Bloch-Kato subgroup $H^1_f(F, \epsilon_X V_s(s+1))$ (*cf.* [28, Thm. 3.1 (ii)] and [29]). In terms of the interpretation of the local cohomology as a group of extension classes, the elements of the subgroup correspond to crystalline extensions.

In view of the Bloch-Kato logarithm map and de Rham Poincaré duality, we have a canonical isomorphism

$$\log : H^1_f(F_v, \epsilon_X V_s(s+1)) \simeq (Fil^{s+1}\epsilon_X H^{2s+1}_{dR}(X_{r_1,r_2}/F_v)(s))^\vee \tag{3.12}$$

(*cf.* [2, §3.4]). Here $Fil^\bullet$ is the Hodge filtration on $\epsilon_X H^{2r+1}_{dR}(X_r/F_v)(r)$ and $(Fil^{r+1}\epsilon_X H^{2r+1}_{dR}(X_r/F_v)(r))^\vee$ the dual arising from the Poincaré pairing.

The composition with the étale Abel-Jacobi map gives rise to the p-adic Abel-Jacobi map

$$AJ_F : CH^{s+1}(X_{r_1,r_2} \otimes F)_{0,\mathbf{Q}} \to (Fil^{s+1}\epsilon_X H^{2s+1}_{dR}(X_{r_1,r_2} \otimes F_v)(s))^\vee. \tag{3.13}$$

There does not seem to be a general conjecture regarding the image and kernel of the p-adic Abel-Jacobi map. However,

(BBK) the Bloch-Beilinson and the Bloch-Kato conjectures suggest to investigate contexts where the $\mathbf{Q}_p$-linearisation $AJ_F \otimes \mathbf{Q}_p$ is an isomorphism or generically so (*cf.* [28, (2.1)]).

In general, the $\mathbf{Q}_p$-linearisation $AJ_F \otimes \mathbf{Q}_p$ need not be injective or surjective. For a relation of the non-triviality of the p-adic Abel-Jacobi map to coniveau filtration and refined Bloch-Beilinson conjecture, we refer to [4, §2].

For later purposes, we recall an explicit description of the middle step $Fil^{s+1}\epsilon_X H^{2s+1}_{dR}(X_{r_1,r_2}/F_v)(s)$ of the Hodge filtration.

For a positive integer $k \geq 2$, let $S_k(\Gamma_1(N), F_v)$ denote the space of cusp form of level $\Gamma_1(N)$ with coefficients in F_v. We have a canonical isomorphism

$$S_{r_1+2}(\Gamma_1(N), F_v) \otimes \mathrm{Sym}^{r_2} H^1_{dR}(A/F_v) \simeq Fil^{s+1}\epsilon_X H^{2s+1}_{dR}(X_{r_1,r_2}/F_v)(s), \tag{3.14}$$

essentially due to Scholl (*cf.* [2, §2.2]) and it arises from

$$f \otimes \alpha \mapsto \omega_f \wedge \alpha.$$

Here ω_f is the normalised differential associated with $f \in S_{r_1+2}(\Gamma_1(N), F_v)$ defined in [2, Cor 2.3] and $\alpha \in \mathrm{Sym}^{r_2} H^1_{dR}(A/F_v)$.

The symmetric power $\mathrm{Sym}^{r_2} H^1_{dR}(A/F_v)$ in turn has the following basis. Let $\omega_A \in \Omega^1_{A/F_v}$ be a non-zero differential. Let $\eta_A \in \Omega^1_{A/F_v}$ be the corresponding element in [2, (1.4.2)]. It turns out that $\{\omega_A, \eta_A\}$ is a basis of Ω^1_{A/F_v}. Thus, $\left\{\omega_A^j \eta_A^{r_2-j} : 0 \leq j \leq r_2\right\}$ is a basis of $\mathrm{Sym}^{r_2} H^1_{dR}(A/F_v)$.

For $0 \leq j \leq r_2$ and f varying over Hecke eigenforms in $S_{r_1+2}(\Gamma_1(N), F_v)$, the wedge products

$$\left\{\omega_f \wedge \omega_A^j \eta_A^{r_2-j} : f, 0 \leq j \leq r_2\right\}$$

thus form a basis of $Fil^{s+1}\epsilon_X H^{2s+1}_{dR}(X_{r_1,r_2}/F_v)(s)$.

3.2.3. *Conjectures.* We state conjectures regarding the non-triviality of generalised Heegner cycles and the image under the étale and p-adic Abel-Jacobi map over anticyclotomic extensions of an imaginary quadratic extension. These conjectures are essentially a compilation of the conjectures due to Mazur, Bloch-Beilinson and Bloch-Kato. For the later conjectures, we refer to [26] and [5].

Let the notations and hypotheses be as in §3.2.1. In particular, N is a positive integer such that $p \nmid N$. For a positive integer $k \geq 2$, let $S_k(\Gamma_0(N), \epsilon)$ be the space of elliptic cusp forms of weight k, level $\Gamma_0(N)$ and neben-character ϵ. Let $f \in S_{r_1+2}(\Gamma_0(N), \epsilon)$ be a normalised newform. In particular, it is a Hecke eigenform with respect to all Hecke operators. Let $N_\epsilon | N$ be the conductor of ϵ. Let $\mathfrak{N}_\epsilon | \mathfrak{N}$ be the unique ideal of norm N_ϵ. The existence follows from the generalised Heegner hypothesis (Hg).

Let $\mathbf{N} : \mathbf{A}_{\mathbf{Q}}^\times / \mathbf{Q}^\times \to \mathbf{C}^\times$ be the norm Hecke character over $\mathbf{Q}$ given by

$$\mathbf{N}(x) = ||x||.$$

Here $||\cdot||$ denotes the adelic norm. Let $\mathbf{N}_K := \mathbf{N} \circ N_{\mathbf{Q}}^K$ be the norm Hecke character over K for the relative norm $N_{\mathbf{Q}}^K$. For a Hecke character $\lambda : \mathbf{A}_K^\times / K^\times \to \mathbf{C}^\times$ over K, let $\mathfrak{f}_\lambda$ (resp. ϵ_λ) denote its conductor (resp. the restriction $\lambda|_{\mathbf{A}_{\mathbf{Q}}^\times}$) as before.

Let b be a positive integer prime to N. Let $\Sigma_{r_1,r_2}(b, \mathfrak{N}, \epsilon)$ be the set of Hecke characters λ such that:

(C1) λ is of of infinity type (j_1, j_2) with $j_1 + j_2 = r_2$ and $\epsilon_\lambda = \epsilon \mathbf{N}^{r_2}$,
(C2) $\mathfrak{f}_\lambda = b \cdot \mathfrak{N}_\epsilon$ and
(C3) The local root number $\epsilon_q(f, \lambda^{-1}\mathbf{N}_K^{-u}) = 1$, for all finite primes q.

Let $\Sigma^{(2)}_{r_1,r_2}(b, \mathfrak{N}, \epsilon)$ be the subset of $\Sigma_{r_1,r_2}(b, \mathfrak{N}, \epsilon)$ whose elements have infinity type $(r_2 + 1 - j, 1 + j)$

for some $0 \leq j \leq r_2$.

Let $\chi \in \Sigma^{(2)}_{r_1,r_2}(b, \mathfrak{N}, \epsilon)$ with infinity type $(r_2 + 1 - j, 1 + j)$ for some $0 \leq j \leq r_2$.

Let M_f (resp. $M_{\chi^{-1}}$) be the Grothendieck motive associated with f (resp. χ^{-1}). By the Künneth formula, the motive $H^{s+1}(X_{r_1,r_2}) \otimes L$ contains $M_f \otimes M_{\chi^{-1}}$ as a submotive for a sufficiently large number field L. For a minimal choice of L, we refer to [4, §4.2]. Let $\pi_{f,\chi}$ be the projector defining the submotive (*cf.* [31]).

Let S_b be a set of representatives for $\mathrm{Pic}(\mathcal{O}_{bN})$ consisting of ideals prime to N. We now regard χ as an ideal Hecke character. Let H_χ be the abelian extension of H generated by the values of χ on S_b. In particular, the extension H_χ depends on the choice of S_b. Let $v = v_\chi$ be the place above p in H_χ induced via the p-adic embedding ι_p.

Let

$$\Delta_\chi = \sum_{[\mathfrak{a}] \in S_b} \chi^{-1}(\mathfrak{a}) \mathbf{N}(\mathfrak{a}) \Delta_{\varphi_\mathfrak{a}} \in CH^{s+1}(X_{r_1,r_2} \otimes H_\chi)_{0,\mathbf{Q}} \otimes L. \tag{3.15}$$

The cycle depends on the choice of representatives S_b. Moreover, it is defined over the extension H_χ by the definition.

Definition 3.4. The generalised Heegner cycle $\Delta_{f,\chi}$ associated with the pair (f, χ) is given by

$$\Delta_{f,\chi} = \pi_{f,\chi}(\Delta_\chi).$$

The generalised Heegner cycle is independent of the choice of representatives S_b at least up to a cycle in the kernel of the complex Abel-Jacobi map (*cf.* [4, Rem. 4.2.4]).

We consider the non-triviality of the cycles $\Delta_{f,\chi}$, as χ varies.

Under the hypotheses (C1)-(C3), the Rankin-Selberg L-function $L(f, \chi^{-1}\mathbf{N}_K^{-u}, s)$ is self-dual with root number -1. The Bloch-Beilinson conjecture accordingly implies the existence of a non-torsion homologically trivial cycle in the Chow realisation of the motive $M_f \otimes M_{\chi^{-1}\mathbf{N}_K^{-u}}$. Perhaps, a natural candidate is the cycle $\Delta_{f,\chi\mathbf{N}_K^u}$. There exists an algebraic correspondence from X_{r_1} to X_{r_1,r_2} mapping the cycle $\Delta_{f,\chi\mathbf{N}_K^u}$ to the $\Delta_{f,\chi}$ (*cf.* [4, Prop. 4.1.1]). One can thus expect a generic non-triviality of the cycles $\Delta_{f,\chi}$, as χ varies.

Based on Mazur's consideration in the case of weight two (*cf.* [26]), an Iwasawa theoretic family of the cycles arises as follows. We first fix a Hecke character $\eta \in \Sigma^{(2)}_{r_1,r_2}(c, \mathfrak{N}, \epsilon)$, for some c prime to N. Let l be an odd prime unramified in K and prime to cN. Let $H_{cNl^\infty} = \bigcup_{n \geq 0} H_{cNl^n}$ be the ring class field of conductor cNl^∞. Let $K_\infty \subset H_{cNl^\infty}$ be the anticyclotomic $\mathbf{Z}_l$-extension of K. Let $G_n = \mathrm{Gal}(H_{cNl^n}/K)$ and $\Gamma_l = \varprojlim G_n$. Let $\mathfrak{X}_l$ be the subgroup of all characters of finite order of the group $\mathrm{Gal}(K_\infty/K) \simeq \mathbf{Z}_l$. As $\nu \in \mathfrak{X}_l$ varies, we consider the non-triviality of the generalised Heegner cycles $\Delta_{f,\eta\nu}$.

Conjecture A. Let the notation be as above. Let $f \in S_{r_{1+2}}(\Gamma_0(N), \epsilon)$ be a normalised newform and η a Hecke character such that $\eta \in \Sigma^{(2)}_{r_1,r_2}(c, \mathfrak{N}, \epsilon)$, for some c prime to N. Suppose that the hypothesis

(Hg) holds. Then, for all but finitely many $\nu \in \mathfrak{X}_l$ we have

$$\Delta_{f,\eta\nu} \neq 0.$$

We also consider the non-triviality of the generalised Heegner cycles under the étale Abel-Jacobi map.

Recall that p is an odd prime prime to N. Let $M_{f,\acute{e}t}$ (resp. $M_{\chi^{-1},\acute{e}t}$) be the p-adic étale realisation of the motive M_f (resp. $M_{\chi^{-1}}$). We accordingly have

$$AJ_{H_{\eta\nu},\acute{e}t}(\Delta_{f,\chi}) \in H^1(H_{\eta\nu}, \epsilon_X(M_{f,\acute{e}t} \otimes M_{\chi^{-1}})).$$

Conjecture B. Let the notation be as above. Let $f \in S_{r_{1+2}}(\Gamma_0(N), \epsilon)$ be a normalised newform and η a Hecke character such that $\eta \in \Sigma^{(2)}_{r_1,r_2}(c, \mathfrak{N}, \epsilon)$, for some c prime to N. Suppose that the hypotheses (ord) and (Hg) hold. Then, for all but finitely many $\nu \in \mathfrak{X}_l$ we have

$$AJ_{H_{\eta\nu},\acute{e}t}(\Delta_{f,\eta\nu}) \neq 0.$$

We finally consider the non-triviality of the generalised Heegner cycles under the p-adic Abel-Jacobi map.

We first recall that the Bloch-Kato subgroup $H^1_f(H_{\eta\nu,v}, \epsilon_X(M_{f,\acute{e}t} \otimes M_{\chi^{-1}}))$ is one-dimensional over $H_{\eta\nu,v}$. Moreover, under the Bloch-Kato logarithm map the corresponding basis is given by $\omega_f \wedge \omega_A^j \eta_A^{r_2-j}$.

Conjecture C. Let the notation be as above. Let $f \in S_{r_{1+2}}(\Gamma_0(N), \epsilon)$ be a normalised newform and η a Hecke character such that $\eta \in \Sigma^{(2)}_{r_1,r_2}(c, \mathfrak{N}, \epsilon)$, for some c prime to N. Suppose that the hypotheses (ord) and (Hg) hold. Then, for all but finitely many $\nu \in \mathfrak{X}_l$ we have

$$AJ_{H_{\eta\nu}}(\Delta_{f,\eta\nu})(\omega_f \wedge \omega_A^j \eta_A^{r_2-j}) \neq 0.$$

Conjecture C (resp. Conjecture B) evidently implies Conjecture B (resp. Conjecture A).

We remind that Conjecture C supports the Bloch-Beilinson and the Bloch-Kato conjecture (*cf.* (BBK)).

As a refinement of Conjecture C, we can ask for a generic mod p non-vanishing of normalised p-adic Abel-Jacobi image. We do not precisely formulate the refinement. In this article, we instead describe results towards the refinement.

3.3. **Non-triviality II.** In this subsection, we prove the non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles modulo p (*cf.* Conjecture C). As the proofs are quite similar to the ones in §3.1, we skip some details.

Unless otherwise stated, let the notations and hypotheses be as in §3.2. In particular, f is a normalised newform of weight $k \geq 2$, level $\Gamma_0(N)$ and η a Hecke character over K. We consider the

non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles $\Delta_{f,\eta\nu}$ as $\nu \in \mathfrak{X}_l$ varies.

In the residually non CM case, our result towards a refinement of Conjecture C is the following.

Theorem 3.5. *Let the notation be as above. Let $f \in S_{r_1+2}(\Gamma_0(N), \epsilon)$ be a normalised newform and $\eta \in \Sigma_{r_1,r_2}(c, \mathfrak{N}, \epsilon)$ a Hecke character of infinity type $(r_2+1-j, 1+j)$, for some c prime to N and $0 \leq j \leq r_2$. In addition to the hypotheses (ord) and (Hg) hold, suppose that*

(1). The residual representation $\rho_f|_{G_K}$ mod $\mathfrak{m}_p$ is absolutely irreducible,
(2). $\mathfrak{N}^-$ is square-free and
(3). $p \nmid \prod_{v|c^-} \Delta_{\eta,v}$.

Then, for all but finitely many $\nu \in \mathfrak{X}_l$ (as $n \to \infty$)

$$v_p\left(\frac{\mathcal{E}_p(f, \eta\nu\mathbf{N}_K^u)}{j!} AJ_{H_{\eta\nu}}(\Delta_{f,\eta\nu})(\omega_f \wedge \omega_A^j \eta_A^{r-j})\right) = 0,$$

where $\mathcal{E}_p(f, \eta\nu\mathbf{N}_K^u) = 1 - (\eta\nu\mathbf{N}_K^u)^{-1}(\overline{\mathfrak{p}})\mathbf{a}_p(f) + (\eta\nu\mathbf{N}_K^u)^{-2}(\overline{\mathfrak{p}})\epsilon(p)p^{r_2+1}$.

Proof. In view of [4, Prop. 4.1.2], it suffices to restrict to the case $r_1 = r_2 = r$.

Let $f^{(p)}$ be the p-depletion of the f as in (3.1). The pair $(d^{-1-j}(f^{(p)}), \eta\mathbf{N}_K^{-1-j})$ satisfies the hypothesis (T) and the toric periods are well defined.

The toric period is given by

$$P_{d^{-1-j}(f^{(p)}),\eta\mathbf{N}_K^{-1-j}}(\nu, n) = \frac{\mathcal{E}_p(f, \eta\nu)}{j!} \sum_{[\mathfrak{a}] \in \Gamma_n} (\eta\nu)^{-1}(\mathfrak{a})\mathbf{N}_K(\mathfrak{a}) \cdot AJ_F(\Delta_{\varphi_\mathfrak{a}\varphi_0})(\omega_f \wedge \omega_A^j \eta_A^{r-j}). \tag{3.16}$$

This is again due to Bertolini-Darmon-Prasanna and the proof is similar to that of Lemma 3.2.

For the non-triviality of these toric periods modulo p, we again first find a pair whose toric periods are congruent modulo p to the periods of interest and such that Theorem 2.1 can be applied to the new pair.

We choose the pair to be $(d^{m(p-2)p-j}(f), \eta\mathbf{N}_K^{m(p-2)p-j})$, where m is a positive integer such that $p-1$ divides $m(p-2)p+1$ and $k+2m(p-2)p-2j \geq 2$. As in the proof of Theorem 3.3, we have

$$P_{d^{m(p-2)p-j}(f),\eta\mathbf{N}_K^{m(p-2)p-j}}(\phi, n) \equiv P_{d^{-1-j}(f^{(p)}),\eta\mathbf{N}_K^{-1-j}}(\phi, n) \bmod \mathfrak{m}_p. \tag{3.17}$$

The non-triviality hypothesis (H) can be verified for the pair in the way as in the proof of Theorem 3.3.

□

Remark. (1). In view of [4, §2.3], it follows that the generalised Heegner cycles are non-trivial in the top graded piece of the coniveau filtration on the middle-codimensional Chow group of X_{r_1,r_2} over the $\mathbf{Z}_l$-anticyclotomic extension of K. The non-triviality is thus an evidence for the refined

Bloch-Beilinson conjecture (*cf.* [4, §2]).

(2). In [21], Howard proved an analogous non-triviality of the étale Abel-Jacobi image of classical Heegner cycles for infinitely many $\nu \in \mathfrak{X}_l$ (*cf.* Conjecture B). The approach is a cohomological adoption of Cornut's approach (*cf.* [11]).

(3). It seems likely that the above approach would also work for the case when f has CM over K.

(4). An analogous non-triviality in the case $l = p$ follows from the results in [10] and [23] (*cf.* [9, §4.3]).

4. p-indivisibility of Heegner points

In this section, we prove the p-indivisibility of the Heegner points over the $\mathbf{Z}_l$-anticyclotomic extension of K when the weight two Hecke-eigen cuspform is p-ordinary. This is a rather straightforward consequence of Theorem A.

Let the notations and hypotheses be as in the introduction. In particular, $\mathfrak{p}$ is a prime above p in K induced by the p-adic embedding ι_p. Let $\mathfrak{p}_n$ be a prime above $\mathfrak{p}$ in K_{cNl^n} induced by ι_p. Let $K_n/\mathbf{Q}_p$ denote the p-adic local field $K_{cNl^n,\mathfrak{p}_n}$ and O_n the integer ring. By defintion, $K_n/\mathbf{Q}_p$ is an unramified extension.

Let f be a p-ordinary Hecke-eigen cuspform as in the introduction. It follows that the corresponding abelian variety B_f is p-ordinary and extends as an abelian scheme over O_n.

Our result regarding the p-indivisibility of the Heegner points is as follows.

Theorem 4.1. *Let the notation be as above. Suppose that f is p-ordinary, (ord) and (Hg) holds. Then, there exists a positive integer n_0 such that for all integers $n \geq n_0$, the Heegner points $\Phi_f(\Delta_{cl^n})$ are p-indivisible in the Mordell-Weil group $B_f(K_n)$.*

Proof. As B_f is p-ordinary and $K_n/\mathbf{Q}_p$ is unramified, it follows that for a non-torsion $P \in B_f(K_n)$, we have $v_p(\log_{\omega_{B_f}}(P)) \geq 1$.

In view of Theorem A, there exists an integer n_0 such that for all integers $n \geq n_0$, the Heegner points $\Phi_f(\Delta_{cl^n})$ are non-torsion. This finishes the proof.

□

Remark. In [16], arithmetic applications of the p-indivisibility of the Heegner point defined over the imaginary quadratic extension K are considered under different hypothesis including the prime l being inert in K. It seems likely that Theorem 4.1 also has similar arithmetic applications.

References

[1] E. Aflalo and J. Nekovář, *Non-triviality of CM points in ring class field towers*, With an appendix by Christophe Cornut. Israel J. Math. 175 (2010), 225–284.

[2] M. Bertolini, H. Darmon and K. Prasanna, *Generalised Heegner cycles and p-adic Rankin L-series*, Duke Math. J. 162 (2013), no. 6, 1033–1148.

[3] M. Bertolini, H. Darmon and K. Prasanna, *Chow-Heegner points on CM elliptic curves and values of p-adic L-functions*, Pacific Journal of Mathematics, Vol. 260, No. 2, 2012. 261-303.

[4] M. Bertolini, H. Darmon and K. Prasanna, *p-adic L-functions and the coniveau filtration on Chow groups*, preprint, 2013, available at "http://www.math.mcgill.ca/darmon/pub/pub.html".

[5] S. Bloch and K. Kato, *L-functions and Tamagawa numbers of motives*, The Grothendieck Festschrift, Vol. I, 333–400, Progr. Math., 86, Birkhäuser Boston, Boston, MA, 1990.

[6] M. Brakocevic, *Anticyclotomic p-adic L-function of central critical Rankin-Selberg L-value*, IMRN, Vol. 2011, No. 21, (2011), 4967-5018.

[7] M. Brakocevic, *Non-vanishing modulo p of central critical Rankin-Selberg L-values with anticyclotomic twists*, preprint, 2011, available at "http://www.math.mcgill.ca/~brakocevic".

[8] A. Burungale, *On the non-triviality of the p-adic Abel-Jacobi image of generalised Heegner cycles modulo p, II: Shimura curves*, preprint, to appear in J. Inst. Math. Jussieu, available at "http://arxiv.org/abs/1504.02342".

[9] A. Burungale, *Non-triviality of generalised Heegner cycles over anticyclotomic extensions: a survey*, preprint, submitted to proceedings of 'p-adic aspects of modular forms', 2014.

[10] F. Castella, *On the exception specialisation of big Heegner points*, preprint 2014, avaialble at "http://www.math.ucla.edu/~castella/".

[11] C. Cornut, *Mazur's conjecture on higher Heegner points*, Invent. Math. 148 (2002), no. 3, 495-523.

[12] C. Cornut and V. Vatsal, *Nontriviality of Rankin-Selberg L-functions and CM points*, L-functions and Galois representations, 121–186, London Math. Soc. Lecture Note Ser., 320, Cambridge Univ. Press, Cambridge, 2007.

[13] C.-L. Chai, *Every ordinary symplectic isogeny class in positive characteristic is dense in the moduli*, Invent. Math. 121 (1995), 439-479.

[14] C.-L. Chai, *Families of ordinary abelian varieties: canonical coordinates, p-adic monodromy, Tate-linear subvarieties and Hecke orbits*, preprint, 2003.

[15] C.-L. Chai, *Hecke orbits as Shimura varieties in positive characteristic*, International Congress of Mathematicians. Vol. II, 295-312, Eur. Math. Soc., Zurich, 2006.

[16] B. Gross and J. Parson, *On the local divisibility of Heegner points*, Number theory, analysis and geometry, 215–241, Springer, New York, 2012.

[17] H. Hida, *Non-vanishing modulo p of Hecke L-values*, In "Geometric Aspects of Dwork Theory" (A. Adolphson, F. Baldassarri, P. Berthelot, N. Katz and F. Loeser, eds.), Walter de Gruyter, Berlin, 2004, 735-784.

[18] H. Hida, *p-Adic Automorphic Forms on Shimura Varieties*, Springer Monogr. in Math., Springer-Verlag, New York, 2004.

[19] H. Hida, *Non-vanishing modulo p of Hecke L-values and applications*, London Mathematical Society Lecture Note, Series 320 (2007), 207-269.

[20] H. Hida, *Elliptic Curves and Arithmetic Invariants*, Springer Monogr. in Math., 2013.

[21] B. Howard, *Special cohomology classes for modular Galois representations*, J. Number Theory 117 (2006), no. 2, 406–438.

[22] M.-L. Hsieh, *On the non-vanishing of Hecke L-values modulo p*, American Journal of Mathematics, 134 (2012), no. 6, 1503-1539.

[23] M.-L. Hsieh, *Special values of anticyclotomic Rankin-Selberg L-functions*, Documenta Mathematica, 19 (2014), 709-767.

[24] N. M. Katz, *p-adic L-functions for CM fields*, Invent. Math., 49(1978), no. 3, 199-297.

[25] Y. Liu, S. Zhang and W. Zhang, *On p-adic Waldspurger formula*, preprint, 2013, available at http://www.math.mit.edu/~ liuyf/ .

[26] B. Mazur, *Modular curves and arithmetic*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983), 185–211, PWN, Warsaw, 1984.

[27] J. Nekovář, *Kolyvagin's method for Chow groups of Kuga-Sato varieties*, Invent. Math. 107 (1992), no. 1, 99–125.

[28] J. Nekovář, *p-adic Abel-Jacobi maps and p-adic heights*, The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), 367–379, CRM Proc. Lecture Notes, 24, Amer. Math. Soc., Providence, RI, 2000.

[29] T. Saito, *Weight-monodromy conjecture for l-adic representations associated to modular forms*, The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), 427–431, NATO Sci. Ser. C Math. Phys. Sci., 548, Kluwer Acad. Publ., Dordrecht, 2000.

[30] C. Schoen, *Complex multiplication cycles on elliptic modular threefolds*, Duke Math. J. 53 (1986), no. 3, 771–794.

[31] A. Scholl, *Motives for modular forms*, Invent. Math. 100 (1990), no. 2, 419–430.
[32] J.-L. Waldspurger, *Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie*, Compositio Math. 54 (1985), no. 2, 173–242.
[33] V. Vatsal, *Uniform distribution of Heegner points*, Invent. Math. 148, 1-48 (2002).
[34] V. Vatsal, *Special values of anticyclotomic L-functions*, Duke Math J., 116, pp 219–261 (2003).
[35] V. Vatsal, *Special values of L-functions modulo p*, International Congress of Mathematicians. Vol. II, 501–514, Eur. Math. Soc., Zürich, 2006.

Department of Mathematics, UCLA, Los Angeles, CA 90095-1555, USA
E-mail address: ashayburungale@gmail.com